

Effect of Four *Trichoderma* Species on Growth and Development of *Rhizoctonia solani*, Isolated From Potato (cv. Spunta), Infected With Black Scurf Disease.

Muna N. Ekrim El-Taher A. Abuhligha Alsadek M. Ghazala

Abeer Masoud Al-Qebrawi Amina Abdul Salam Al-Qamoudi

University of Tripoli, Faculty of Agriculture

Abstract:

Rhizoctonia solani isolates were obtained from infected potato tubers (cv. Spunta). Four isolates of *Trichoderma* spp., namely *Trichoderma harzianum*, *T. viride*, *T. longibrachiatum*, and *T. album*, were applied to test their antagonistic effects on growth and development of the obtained isolates of *R. solani*. Interactions between the antagonistic fungal isolates and *R. solani* were studied in dual culture, antagonism and volatile, nonvolatile techniques. All the tested *Trichoderma* isolates showed significant antagonistic effects on the tested *R. solani* isolates. They produced inhibition zones in front of the *R. solani* colonies, but in varying degrees. *T. harzianum* isolates were able to overgrow the mycelium of *R. solani*, compared to those of the other *Trichoderma* isolates. All the antagonists suppressed the formation of sclerotia. This resulted in the reduction of the mycelial growth of the *R. solani* isolates. The results implied that the extent of inhibition by *T. harzianum* provides the use of excellent potential antagonists capable of controlling the pathogenicity of *R. solani* on Potato.

Key words: Potato, fungal bioagents, antagonism, parasitism, *Rhizoctonia solani*

Introduction:

Potato (*Solanum tuberosum* subsp. *Tuberosum* L), which belongs to family Solanaceae, is considered one of the most important crops in the world after rice, barley, wheat and maize (Ahmad *et al.*, 1995 and Rauscher *et al.*, 2006). Moreover, potato is one of the most important agricultural and food crops in Libya, where it is grown throughout the year in two seasons, i.e. autumn and spring. It is predicted that by 2050 the world overall population will approximately reach 9 billion. Therefore, to feed this increasing world population, a raise of about 70% in agriculture food production is necessary (Raney T, 2009).

Biotic stress factors involving fungi, bacteria, virus, nematodes, weeds, and insects cause a yield loss up to 31–42 % (Moustafa-Farag *et al.*, 2020). Among these factors, fungal pathogens are considered the most severe limiting factor for crop production worldwide. Greater than 10,000 spp. of fungi are considered responsible for plant disease incidence.

Biological control mechanisms are considered as significant measures for disease management because chemical fungicides adversely affect other non-target organisms (Kohl *et al.*, 2019). There are many organisms that infect potato crop, e. g. *Rhizoctonia solani*, which is the subject of the present study, the causative agent of potato black scurf and stem canker (Raju *et al.*, 2003, Viterbo *et al.*, 2006, Lace *et al.*, 2015, and Shukla, *et al.*, 2012).

There were many evidences that support the fact that some microorganisms can cause growth inhibition of pathogenic species by decreasing their metabolism and/or establishing a parasitic relationship (Panth *et al.*, 2020). Additionally, the application of biological control agents (BCAs) with reduced concentrations of chemicals can stimulate disease suppression in a similar manner as high doses of chemical fungicide treatment (Hyder *et al.*, 2017). Around 90% of fungal biocontrol agents, which are applied, against pathogenic microorganisms, are belonged to different strains of *Trichoderma* (Hermosa *et al.*, 2012). *Trichoderma* was isolated for the first time in 1794 from soil and decomposing organic matter (Persoon,

1794). Throughout the world, currently greater than 60% effective bio-fungicides were obtained from *Trichoderma* (Abbey *et al.*, 1919). For example, in India, approximately 250 *Trichoderma*-derived bio fungicides products are employed, however, Indian farmers are still relying on synthetic chemical fungicides to a great extent (Singh *et al.*, 2009). Different strains of *Trichoderma* (teleomorph *Hypocrea*) are belonging to fungi imperfecti, which lacking sexual stages in their life cycles (Van Wees *et al.*, 2008). These fungi are rapid colonizers, invasive, filamentous, opportunistic, avirulent and exhibit a symbiotic relationship with plants. In pathogen-contaminated soils, they are not only improve plant growth but also inhibit pathogen growth through several antagonistic mechanisms (Vinale *et al.*, 2008 and Lorito *et al.*, 2010). *Trichoderma* exhibits antagonistic behavior against several phytopathogenic organisms, including bacteria, nematodes and especially fungi, by inhibiting their growth either by direct interaction (e.g., hyperparasitism, competition for nutrient and space, and by antibiosis (Zhang *et al.*, 2017), or indirectly by improving plant growth, and enhancing stress tolerance, active uptake of nutrients and bioremediation of contaminated rhizosphere, as well as providing plants with several secondary metabolites, enzymes and pathogen proteins (Kumar, 2013).

Therefore, the current work aims to investigate the antagonistic effect of four of *Trichoderma* species, e.g., *T. viride*, *T. harzianum*, *T. brachiatum*, and *T. album*, on five isolates of *Rhizoctonia solani*, isolated from infected potato tubers.

Materials and methods:

Sources of pathogen and bioagents isolates :

Infected potato (cv. Spunta) samples were collected randomly from popular markets. The antagonistic fungi *Trichoderma viride*, *T. harzianum*, *T. longibrachiatum*, and *T. album*, applied throughout the present study, were obtained from Botany Department at Saba Baha Agriculture Faculty, Alexandria University.

Isolation of the pathogen:

Potato tubers infected with black scurf disease were surface sterilized by 0.2% sodium hypochlorite (NaOCl) for two minutes and extensively washed with sterile distilled water. Clot like black spot on

the surface of potato were peeled and placed on petri dishes containing potato dextrose agar medium (PDA), and then incubated at 25 ± 2 °C for 5-7 days. The fungal culture growth was purified by subculturing and identified by light microscope after staining with trypan blue dye. Pure culture of *R. solani* was stored at 5 °C in tubes and petri plates containing PDA and sealed with parafilm (Juan-Abgona *et al.*, 1996).

Dual culture method:

In vitro antagonistic activity of *T. viride*, *T. harzianum*, *T. longibrachium* and *T. album* against *Rhizoctonia solani* isolates was studied in dual culture technique by following the method by Kucuk and Kivanc (2003). Petri dishes (90 mm) containing 20 ml of sterile PDA were inoculated with a 5 mm in diameter plug of 7- day- old pure culture of antagonistic fungi and pathogens. One mycelial disc of each fungus was placed on opposite poles of PDA plates and incubated at 25°C in incubator and radial growth of pathogen was measured at 7 days intervals. Control petri dishes were inoculated with pathogens and a sterile agar plug. Linear growth of the tested fungi was measured when pathogenic fungi have completely covered the surface of the medium in the control treatment. Five replicates were maintained for each treatment. The percentage of inhibition was calculated using the following formula: Percentage of inhibition = $(A1 - A2) / A1 \times 100$, Where A1 = Area covered by the pathogen in control, A2 = Area covered by pathogen in dual culture.

Ranking based on Bell *et al.* (1982) scale was used in order to evaluate the antagonistic potential of isolates of *Trichoderma* interactions between *Rhizoctonia solani* isolates (Table 1).

Table 1. Scaled to the classes for antagonism of *Trichoderma* spp. with *R. solani* isolates (adapted from Bellet *al.*, 1982)

Class	Characteristics
1	<i>Trichoderma</i> grows and covers completely all colonies of <i>R. solani</i> and medium surface.
2	<i>Trichoderma</i> grows and covers 2/3 of medium surface.
3	Antagonists and phytopathogen colonize each one, half of the medium surface and no one seems to dominate the other.
4	<i>R. solani</i> colonizes 2/3 of medium surface.
5	<i>R. solani</i> grows and covers completely all colonies of <i>Trichoderma</i> and the medium surface.

Effect of volatile substance produced by *Trichoderma* spp on growth of the pathogen:

The method described by Dennis and Webster (1971) was adopted. Petriplates containing 20 ml of PDA were inoculated separately with 5 mm disc of antagonists and incubated for 5 hours. After this lid of each plate was replaced by a bottom containing PDA previously inoculated with the disc of the pathogen and sealed together with paraffin film. The control sets did not contain the antagonist. The cultures were incubated at 25°C. The studies were carried out in three replicates. Radial growth was measured at 3 and 5 day intervals and percent inhibition was determined using the formula: Percent inhibition = $C2 - C1 / C2 \times 100$. Where, C2 means growth of *R. solani* in control and C1 means growth of *R. solani* in treatment.

Non-volatile activity

Non-volatile activities of fungal biocontrol agents were tested as described by Jariwala *et al.*, (1991), 1 ml of spore suspension (1×10^5 cfu / ml) was inoculated in Potato Dextrose (PD) broth and incubated at room temperature for 1 week. The fungal mat and the spores were removed by filtration through double layer filter paper, followed by

centrifugation at 3000rpm for 15min. The supernatant was used for antibiotic activity. Culture filtrates were added to PD Agar medium at 25%, 50%, 75% and 100% concentration, the pH was adjusted to 6.8 ± 0.2 . Medium was then sterilized and poured in the sterile Petri plates. Five day old actively growing *R. solani* cultures were removed from the edge of the colony using 5mm diameter cork borer and placed at the center of these culture medium for each isolates and the plates were incubated 25 ± 2 °C. Five replicates were maintained for each concentration. Plates containing PDA medium with pathogens alone served as control. Radial growth of the fungal colony was measured 7 days after incubation. Percentage of inhibition was calculated using the formula described earlier.

Experimental design and statistical analysis in *vitro* studies were done using a completely randomized design (CRD), each treatment with five replicates. Data were analyzed statistically using the (Minitab) Computer Program. The means were compared using the Fisher LSD ($P = 0.05\%$) (Welkowitz *et al.*, 1982).

Experimental results:

In dual culture assays, according to results of Table 2, significant inhibition ratios of *R. solani* hyphal growth was obtained in all *Trichoderma* treatments. However, inhibition ratios differed according to the species of *Trichoderma* and the tested isolates of *Rhizoctonia* isolate. *T. viride* gave the highest inhibition ratio (82.96%) on isolate No. 1 of *R. solani*, whereas the lowest inhibition by *T. viride* was obtained on isolate No. 5. On the other hand, the highest inhibition ratio (79.63%) was obtained by *T. harzianum* isolate No.1 of *Rhizoctonia*. The least inhibition ratio (68.52%) was obtained in *Rhizoctonia* isolate No.5. Moreover, the highest inhibition ratio (72.96%) was detected by *T. longibrachiatum*, and was recorded by *Rhizoctonia* isolate No.3, whereas the lowest ratio (54.07%) was detected in *Rhizoctonia* isolate No. 2. Finally, *T. album* treatment showed highest inhibition value in isolate No.1 treatment, and the lowest in isolate No. 4. Furthermore, the *T. harzianum* showed the highest mean of inhibition (73.6%), with the highest levels against isolate No. 1 (61.98 %). Based on the above, all the antagonists were


suppressed sclerotia formation on antagonized portion of *Rhizoctonia* isolates hyphal after 7 days of inoculation.

Table 2. Effect of *Trichoderma* spp on mycelial growth inhibition of *Rhizoctonia solani* isolates.

Bioagents (B)	Growth inhibition %					Mean
	Isolates of <i>Rhizoctonia solani</i> (I)					
	1	2	3	4	5	
<i>T. viride</i>	82.96	72.08	70.74	68.52	63.07	71.4
<i>T. harzianum</i>	79.63	74.85	75.18	70.00	68.52	73.6
<i>T. longibrachiatum</i>	72.59	54.07	72.96	65.56	66.33	66.30
<i>T. album</i>	74.74	60.74	55.22	52.92	53.33	59.39
Control	0.00	0.00	0.00	0.00	0.00	0.00
Mean	61.98	52.35	54.82	51.40	50.25	54.16
LSD at 0.05 (B) = 0.02 LSD0.05 (I) =0.02 LSD0.05 (B x I) =0.21						

Antagonistic potential of *Trichoderma* spp. against *Rhizoctonia solani* in vitro of the four antagonist *Trichoderma* spp evaluated for potential biocontrol of *R. solani*, only in isolates 1, 2, 3, and 4, showing satisfactory action depending on class 1 of antagonism, adapted from Bell *et al.* (1982). The other isolates of *Trichoderma* spp. showed mild antagonistic potential, pertaining to class 2 of antagonism. In contrast to that, *T. viride* in isolate 5 didn't show high antagonistic action against *Rhizoctonia solani* (Fig.1), whereas isolate T4 (*T. album*) in isolate 5 didn't show antagonistic potential against the pathogen pertaining to class 3. This could be due to the preservation of the isolate, which was obtained from the Micoteca URM collections. According to Smith and Onions (1994), the preservation of microorganisms may alter or inactivate important

physiological characteristics such as pathogenicity, sporulation or enzymes liberation and other compounds. Besides mycelial growth, the level of sporulation of antagonist isolate was also observed. The production of spores of *T.viride* isolates 1, 2, and 3, *T.harzianum*, isolate 3 presented hardly produced spores while a slight decrease in sporulation production in *T.album*, isolate 5, on the other hand *T. longibrachiatum* was not affected when in contact with pathogen, . Batista (2002) reported that the process of sporulation would be possibly a favorable characteristic to antagonists, therefore, new inoculum was desirable in the presence of pathogen, inhibiting its actions, due to greater density of antagonist inoculum.




Isolate 5				
	<i>T. viride</i>	<i>T. harzianum</i>	<i>T. longibrachiatum</i>	<i>T. album</i>

Fig 1. The antagonistic effect of *Trichoderma* spp against *Rhizoctonia solani* isolates

Effect of volatile substance produced by *Trichoderma* spp. on growth of the pathogen.

Determination of inhibition percentages was carried out in two time periods i.e. three days and five days after inoculation. According to the obtained results of the first period, inhibition percentages significantly differed according to the tested *R. solani* isolate and the applied bioagent. In *T. viride* treatment, the highest inhibition % was recorded in *R. solani* isolate 2 (87.50%), whereas the least inhibition (37.12%) was detected in isolate 5. Similarly, in *T. harzianum* treatment, the highest inhibition (68.78%) was detected in isolate 5 of *R. solani*, and the least percent was obtained on isolate 4 (40.67%). The highest inhibition% in *T. longibrachiatum* was similarly detected in isolate 5 (63.52%), whereas isolate 1 realized the least percent (37.04%). In contrast to that, *T. album* showed the highest inhibition% on isolate 2 (60.25%) and the least on isolate 5 (28.15%). In the second tested time period (five days of inoculation), the highest inhibition values were produced by *T. harzianum* on isolate 5 (70.90%), *T. viride* on isolate 2 (64.68%), *T. album* on isolate 2 (57.31%). On the other hand, the lowest inhibition values were produced by *T. viride* on isolate 5 (6.04%), *T. longibrachiatum* on

isolate 2 (11.95), *T. album* on isolate 1 (29.44%), and *T.harzianum* on isolate 3 (31.11%) (Table 3).

Table 3. Volatile activity of *Trichoderma* spp. against *R. solani* isolates in two time periods.

Bioagents(B)	Growth inhibition (%) 3days after inoculation					Me an	Growth inhibition (%) 5 days after inoculation					Me an
	Isolates of <i>Rhizoctonia solani</i> (I)						Isolates of <i>Rhizoctonia solani</i> (I)					
	1	2	3	4	5		1	2	3	4	5	
<i>T. viride</i>	40.73	87.50	47.00	40.00	37.12	50.47	37.88	64.68	24.44	24.97	6.04	31.60
<i>T. harzianum</i>	43.07	70.17	60.00	40.67	78.78	58.54	32.77	50.00	31.11	31.85	70.90	43.33
<i>T. longibrachiatum</i>	37.04	44.17	44.33	37.33	63.52	45.28	13.84	11.95	23.33	21.38	32.54	20.61
<i>T. album</i>	35.04	60.25	38.00	36.67	28.15	39.62	29.44	57.31	32.22	32.25	23.08	34.86
Control	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Mean	31.17	52.42	37.87	30.93	41.51	38.78	22.79	36.79	22.22	22.09	26.51	26.08
	LSD _{0.05} I, B,T = 0.57 LSD _{0.05} IxB= 0.23 LSD _{0.05} IxT, BxT = 0.15 LSD _{0.05} IxBxT = 0.33											

Non-volatile activity

Effect of different concentrations of *Trichoderma* spp. (25%, 50%, 75% and 100%) on mycelial growth of the tested isolates of *R. solani* were tested and results were demonstrated in Table. 4, Where:

(1) ***T. viride* treatment:** the highest inhibition values (25%, 50%), were detected in isolates 3 (25%), isolate 1, (100%), and 100% inhibition% in all isolates at conc, 75%, and 100%. The lowest inhibition levels at conc. 25 and 50% were obtained in isolates 3 (25%) and isolate 1 (100%), respectively. At concentrations 75 and 100%, inhibition % was 100% for all the tested isolates.

(2) ***T. harzianum* treatment:** The highest inhibition values at concentrations 25, 50, and 75% were obtained in isolates 3 (91.35%), isolate 2 (25%), and isolate 1 (100), respectively. At conc. 100%, inhibition% was 100% for all the tested isolates. The lowest inhibition values at conc. 25, 50 and 75% were detected in isolate 1 (0.00%) and isolate 4 (21.67%), respectively. At concentration 100%, inhibition % were 100 for all the tested isolates.

(3) ***T. longibrachiatum* treatment:** The highest inhibition values at concentrations 25, 50, and 75%, and 100% were obtained in isolate 2 (11.33%), isolate 2 (46.39%), isolate 1 (81.94%), and isolates 2 and 4 (100%). The lowest inhibition values at concs. 25, 50, 75%, and 100% were detected in isolate 1 (0.00%), isolate 1 (0.00%), isolate 3 (16.25%), and isolate 3 (60.63%), respectively.

(4) ***T. album* treatment:** The highest inhibition values at concs. 25%, 50%, 75%, and 100% were detected in isolate 3 (11.41%), isolate 2 (54.34%), isolate 2 (79.13%), isolates 1,2 (100%). The lowest inhibition values at concs. 25, 50, and 75%, 100% were isolates 1,2 ,4 (0.00), isolate 1 (0.00), isolate 4 (10.24%), and in isolate 5 (47.12%).

Table 4. Effect of different concentrations of the tested *Trichoderma* spp on mycelial growth of *Rhizoctonia solani* isolates.

Treatements		Growth inhibition (%)					Mean
Bioagent (B)	Concentration % (C)	Isolates of <i>Rhizoctonia solani</i> (I)					
		1	2	3	4	5	
<i>T. viride</i>	25	0.00	8.23	25.00	7.69	3.73	8.93
	50	100.00	10.70	49.50	13.75	5.59	35.908
	75	100.00	100.00	100.00	100.00	100.00	100.00
	100	100.00	100.00	100.00	100.00	100.00	100.00
Control	-	0.00	0.00	0.00	0.00	0.00	0.00
Mean		60.00	43.79	54.90	44.29	41.86	88.967
<i>T. harzianum</i>		1	2	3	4	5	Mean
	25	0.00	9.21	13.00	2.14	3.33	5.536
	50	0.00	25.00	31.00	3.93	12.78	14.542
	75	100.00	77.09	46.00	21.67	22.22	53.396
	100	100.00	100.00	100.00	100.00	100.00	100.00
Control	-	0.00	0.00	0.00	0.00	0.00	0.00
Mean		40.00	42.26	38.00	25.55	27.67	34.694
<i>T. longibrachiatum</i>		1	2	3	4	5	Mean
	25	0.00	11.33	4.38	2.80	1.42	3.986
	50	0.00	46.39	11.41	10.49	19.33	17.524
	75	81.94	74.18	16.25	20.37	22.67	26.694
	100	88.79	100.00	60.63	100.00	65.28	82.94
Control	-	0.00	0.00	0.00	0.00	0.00	0.00

Mean		34.15	46.38	18.53	26.73	21.74	26.228
<i>T. album</i>		1	2	3	4	5	Mean
	25	0.00	0.00	11.41	0.00	2.82	2.846
	50	0.00	54.34	41.30	6.79	7.54	21.199
	75	74.18	79.13	51.00	10.24	21.14	42.91
	100	100.00	100.00	59.00	53.89	47.12	71.824
Control	-	0.00	0.00	0.00	0.00	0.00	0.00
Mean		34.84	46.69	32.54	14.18	15.72	27.755
$LSD_{0.05} I = 0.465$ $LSD_{0.05} CXI = 1.406$ $LSD_{0.05} C = 0.465$ $LSD_{0.05} B = 0.416$ $LSD_{0.05} BXCXI = 2.081$							

Discussion:

R. solani is widespread and responsible for severe damage to many worldwide economically important agricultural and horticultural crops (Grosch *et al.*, 2006). As for potato high yield losses were reported reaching up to 20 % (Grosch *et al.*, 2005). Strategies to control *Rhizoctonia* diseases are limited because of its ecological behavior, large host range and the high survival rate of sclerotia under various environmental conditions (Groth & Bond, 2006). A lot of fungal biota have been reported to be effective biocontrol agents of *R. solani* on potato. Among these are species of *Trichoderma* (Brewer & Larkin, 2005; Grosch *et al.*, 2006 and Wilson *et al.*, 2008). A biocontrol agent may act against pathogens by using one or more of the following mechanisms: competition, antibiosis, and parasitism as well as activating host defense mechanisms (Papavizas & Lumsden, 1980). All isolates were tested against *R. solani* isolate in dual culture. Colony and hyphal interactions of these fungal isolates with *R. solani* on PDA are listed in the Tables. Results of the antagonism tests showed that *T. harzianum*, *T. viride*, *T. longibrachiatum* and *T. album* species were able to overgrow the mycelium of *R. solani* which is compatible with (Chand and Logan, 1984; Anderson, 1988; Nicoletti *et al.*, 2004). Antagonism is not the same in *Trichoderma* spp. because different strains or isolates of the same species can exhibit varying biocontrol potential against *R. solani*. The strains or isolates which

posses genes are efficiently and rapidly expressed involved in antagonist activity against *R. solani* are better antagonists (Schermer *et al.*, 2009). Until now *Trichoderma spp.* have been successfully applied to diseases of *R. solani* mostly in Greenhouse studies with few studies conducted under field conditions (Da Silva *et al.*, 2012), and this is somewhat consistent with our results.

For the control of *R. solani* their extracellular metabolites have been continuously used as biological fungicides (Eziashi *et al.*, 2007). Moreover, *Trichoderma spp.* more usefull for nutrients as compared to a pathogen which provides them a competitive advantage (Sarrocchio *et al.*, 2017). As far as interaction between *Trichoderma spp.* and *R. solani* is concerned, the mycoparasitism is regarded as a major activity. *Trichoderma spp.* sense small molecules released by *R. solani*. Some of these molecules may be released by proteases enzymes.

Moreover, the *Trichoderma spp.* are showing affinity of cell wall of *Trichoderma spp.*, and *R. solani* which then lead to host cell wall penetration (Schuster & Schmoll, 2010). *Trichoderma spp.* are also suppressing *R. solani* by producing antifungal compounds. Moreover, isolates of *Trichoderma* excrete some growth inhibitory substances. Of these, alkyl pyrones, isonitriles, polyketides, peptaibols diketopiperazines, sesquiterpenes, and steroids have frequently been associated with biocontrol activity (Howell, 1998; Sivasithamparan and Ghisalberti, 1998). On the other hand, Vinale *et al.* (2008) reported that *Trichoderma spp.* produced secondary metabolites such as antibiotics (6-pentyl- α -pyrone (6pp), isocyanide derivatives), acids (heptelidic and konigic acid), peptaibols and cell wall degrading enzymes (CDWE) that are implicated in inhibits of radial growth of many phytopathogenic fungi (Verma *et al.*, 2007). Hyperparasitism and volatile metabolites may be involved in the inhibition of *Fusarium oxysporum*, *Rhizoctonia solani* and *Botrytis cinerea* (Naeimi *et al.*, 2010). Cell wall degrading enzymes (CWDEs) such as chitinase, glucanase, and proteases are thought to be closely related to the mycoparasitism of *Trichoderma* strains (Marzano & Altomare, 2013; Pandey *et al.*, 2005 and Saravanakumar *et al.*, 2016). Inhibitory volatile substances such as alkylpyrones may also contribute to the biocontrol activity of some *Trichoderma* strains (Claydon *et al.*, 1987 and Devaki *et al.*,

1992). The antifungal compounds include antibiotics, mycotoxins and lower weight secondary compounds (Schuster and Schmoll ,2010). *Trichoderma spp.* are also well knowing plant growth regulators. They proliferate root and increase the yield by uptake of nutrients (De França *et al.*, 2015). As compared to fungicides the effect of *Trichoderma spp.* against *R. solani* is higher because it persists in soil for a long period after application (Gajera ,*et.al.*,2016). Activation of biocontrol genes varied with various *Trichoderma spp.* are only where there is contact with *R. solani* (Harman and Kubicek, 2002).

Conclusion

The work indicates that fungal biocontrol agents were effective against *R. solani*. These biocontrol agents inhibit the pathogen by antibiosis (volatile or non-volatile) or by parasitism of these methods.

Acknowledgement

The authors are grateful to Prof. Dr. Ibrahim Abdel-Salam El-Samra and Prof. Dr. Mostafa Abd-El-Azeem Amer at Botany Department at Saba Bacha Agriculture Faculty, Alexandria University. for kindly providing *Trichoderma spp.* for our work and providing the necessary facilities for the smooth conduct of this work We are also thankful to Dr.Redha Aboshagor at Agronomy Department, Faculty of Agriculture, University of Tripoli for his help in statistical analysis.

Reference

1. Abbey JA, Percival D, Abbey L, Asiedu SK, Prithiviraj B and Schilder A(2019). Biofungicides as alternative to synthetic fungicide control of grey mould (*Botrytis cinerea*) prospects and challenges. Biocontrol. Sci. Technol., 29, 207–228.
2. Ahmad I, Soomro MH, Khalid S, Iftikhar S, Munir A and Burney K (1995). Recent distributional trends of potato disease in Pakistan. *Proc. of Nat. Sem.* April, 23-25. PARC, Islamabad. PP 3-17.
3. Anderson HA, Bracewell JM and Fraser AR(1988). 5-Hydroxymaltol and mycophenolic acid, secondary metabolites from *Penicillium echinulatum*. Trans British Mycol Soc 91: 649-651.
4. Batista D C (2002). *Manejo integrado da murcha de Fusário do maracujazeiro*. Ms Sc. Thesis, Universidade Federal Rural de Pernambuco, Recife, PE.. 89pp
5. Bell DK, Wells HD and Markham CR (1982). In vitro antagonism of *Trichoderma* species against six fungal plant pathogens, *Phytopathology*, 72-379-382.
6. Brewer MT and Larkin RP (2005). Efficacy of several potential biocontrol organisms against *Rhizoctonia solani* on potato. *Crop Protect*, 24,939-950,doi:10.1016/J.cropro.
7. Chand T, Logan C (1984). Antagonists and parasites of *Rhizoctonia solani* and their efficacy in reducing stem canker of potato under controlled conditions. Trans British Mycol Soc 83: 107-112.
8. Claydon N, Allan M, Hanson IR, Avont AG (1987). Transactions of the British Mycological Society 88: 503-513.
9. Da Silva JC, Torres DB, Lustosa DC, de Filippi MCC and da Silva GB (2012). Rice sheath blight biocontrol and growth promotion by *Trichoderma* isolates from the Amazon. Amazon Journal of Agricultural and Environmental Sciences. www.ajaes.ufra.edu.br. Accessed on 9/4/2021
10. De França SKS, Cardoso AF, Lustosa DC, Ramos EMLS and De Filippi MCC (2015). Biocontrol of sheath blight by *Trichoderma asperellum* in tropical lowland rice. *Agron Sustain Dev* 35: 317-324.

11. Dennis C and Webster J (1971). Antagonistic properties of species-groups of Trichoderma: II. Production of volatile antibiotics. 57: 41-48.
12. Devaki NS, Bhat SS, Bhat SG, Manjunatha KR (1992). Antagonistic activities of *Trichoderma harzianum* against *Pythium aphanidermatum* and *Pythium myriotylum* on tobacco. Jour Phyto 136: 82-87.
13. Eziashi EI, Omamor IB, Dimaro-Oruade EA and Ogunkanmi LA (2007). Control of phytotoxin from *Ceratocystis paradoxa* using *Trichoderma species* phytotoxins on oil palm (*Elaeis quineensis* Jacq.) sprouted seeds. Plant Pathol J 6: 324-329.
14. Gajera HP, Hirpara DG, Katakpara ZA, Patel SV and Golakiya BA (2016). Molecular evolution and phylogenetic analysis of biocontrol genes acquired from SCoT polymorphism of mycoparasitic *Trichoderma koningii* inhibiting phytopathogen *Rhizoctonia solani* Kuhn. Infect Genet Evol 45: 383-392.
15. Grosch R, Faltin F, Lottmann J, Kofoet A and Berg G (2005). Effectiveness of three antagonistic bacterial isolates to suppress *Rhizoctonia solani* Kuhn on lettuce and potato. Canadian Journal of Microbiology, 51: 345 – 353.
16. Grosch R, Scherwinski K, Lottmann J and Berg G (2006). Fungal antagonists of the plant pathogen *Rhizoctonia solani*: Selection, control efficacy and influence on the indigenous microbial community. Mycological Research, 110: 1464 – 1474.
17. Groth DE and Bond JA (2006). Initiation of rice sheath blight epidemics and effects of application timing of azoxystrobin on disease incidence, severity, yield and milling yields. Plant Dis. 90: 1073–1078.
18. Howell CR (1998). The role of antibiosis in biocontrol. In: *Trichoderma and Gliocladium*, Vol 2, (Eds., G.E, Harman and C.P. Kubicek), Taylor and Francies, London. pp.173-184.
19. Hermosa R, Viterbo A, Chet I and Monte E (2012). Plant-beneficial effects of *Trichoderma* and of its genes. Microbiology, 158: 17–25.
20. Hyder S, Inam-ul-Haq M, Bibi S, Humayun A, Ghuffar Sand Iqbal S (2017). Novel potential of *Trichoderma spp.* as biocontrol agent. J. Entomol. Zool. Stud., 5: 214–222.

21. Jariwala S, Vinaykumari and Bharat R (1991). Antagonistic activity of some fungi against *Alternaria solani* and *Drechlera oryzae*. *Acta Botanica Indica*, 19(2): 217 - 223.
22. Juan-Abgona RV, Katsuno N, Kageyama K and Hyakumachi M (1996). Isolation and identification of hypo virulent *Rhizoctonia* spp. From soil. *Plant Pathol.* 45(5):896-904.
23. Köhl J, Kolnaar R and Ravensberg WJ (2019). Mode of action of microbial biological control agents against plant diseases: Relevance beyond efficacy. *Front. Plant Sci.* 10, 845.
24. Kucuk Cand Kivanc M (2003). Isolation of *Trichoderma* spp. and determination of their antifungal, biochemical and physiological features. *Turkish Journal of Biology* 27: 247 –253.
25. Kumar, S (2013). A biological weapon for managing plant diseases and promoting sustainability. *Int. J. Agric. Sci. Med. Vet.*, 1, 106–121.
26. Lace B, Genre A, Woo S, Faccio A, Lorito M, Bonfante and Gate P (2015). crashing arbuscular mycorrhizas: In vivo imaging shows the extensive colonization of both symbionts by *Trichoderma atroviride*. *Environ. Microbiol. Rep.* 7, 64–77.
27. Lorito M, Woo S L, Harman G E and Monte E (2010). Translational research on *Trichoderma*: From 'omics to the field. *Ann. Rev. Phytopathol.*, 48, 395–417.
28. Marzano M, Gallo A, Altomare C (2013). Improvement of biocontrol efficacy of *Trichoderma harzianum* vs. *Fusarium oxysporum* f. sp. *lycopersici* through UV-induced tolerance to fusaric acid. *Biological Control* 67: 397-408.
29. Moustafa-Farag M, Almoneafy A, Mahmoud A, Elkelish A, Arnao MB, Li L. and Ai S (2020). Melatonin and Its Protective Role against Biotic Stress Impacts on Plants. *Biomolecules*, 10, 54.
30. Naeimi S, Okhovvat SM, Javan-Nikkhah M, Vágvölgyi C, Khosravi, V (2010). Biological control of *Rhizoctonia solani* AG1-1A, the causal agent of rice sheath blight with *Trichoderma* strains. *Phytopathol Mediterr* 49: 287-300.
31. Nicoletti R, De Stefano M and, De Stefano S (2004). Antagonism against *Rhizoctonia solani* and fungitoxic metabolite production by some *Penicillium* isolates. *Mycopathologia* 158: 465-474.

32. Rauscher GM, Smart CD, Simko I, Bonierbale M, Mayton H, Greenland A and Fry W E (2006). Characterization and mapping of RPi-ber, a novel potato late blight resistance gene from *Solanum berthaultii*. *Theor. Appl. Genet.*, 112: 674-687.
33. Pandey KK, Pandey PK, Upadhyay JP (2005). Mycoparasitism of *Trichoderma* spp. on *Fusarium* and *Rhizoctonia*. *J Mycol Plant Pathol* 35: 174-176.
34. PanthM, Hassler SC and Baysal-Gurel F (2020). Methods for mangament of soilborne diseases in crop production 10(1):16. <http://doi-org/10.3390/agriculture10010016>.
35. Papavizas GC and, Lumsden R D (1980). Biological-control of soilborne fungal propagules. *Annu. Rev. Phytopathol* 18: 389-413.
36. Persoon CH (1974). *Disposita methodical fungorum*. Romers. *Neues. Mag. Bot.*, 1, 81–128.
37. Raju NS, Niranjana SR and Shetty HS (2003). Effect of *Pseudomonas fluorescens* and *Trichoderma harzianum* on headmoulds and seed qualitties of Sorghum. *Crop Improv. (India)* , 30, 6–12.
38. Raney T (2009). *The State of Food and Agriculture: Livestock in the Balance*; Food and Agriculture Organization of United Nations: Rome, Italy. http://www.maff.go.jp/primaff/koho/seminar/2010/attach/pdf/100513_02.pdf. 10.4.2021.
39. Saravanakumar K, Yu CJ, Dou K, Wang M, Li Y (2016). Overview of Mechanisms of Biological Control. *Bio Cont* 94: 37-46.
40. Sarrocco S, Matarese F, Baroncelli R, Vannacci G and Seidl-Seiboth V (2017). The constitutive endopoly galacturonase TvPG2 regulates the induction of plant systemic resistance by *Trichoderma virens*. *Phytopathology*. <http://doi.org/10.1094/PHYTO-03-16-0139-R>.
41. Scherm B, Schmoll M, Balmas V, Kubicek C and Migheli Q (2009). Identification of potential marker genes for *Trichoderma harzianum* strains with high antagonistic potential against *Rhizoctonia solani* by a rapid subtraction hybridization approach. *Curr Genet* 55(1): 81-91.

42. Schuster A and Schmoll M (2010). Biology and biotechnology of *Trichoderma*. Appl Microbiol Biotechnol 87: 787-799.
43. Shukla N, Awasthi RP, Rawat L and Kumar J (2012). Biochemical and physiological responses of rice (*Oryza sativa* L.) as influenced by *Trichoderma harzianum* under drought stress. Plant Physiol. Biochem., 54, 78–88.
44. Singh HB, Singh BN, Singh SP, Singh SR and Sarma BK (2009). Biological control of plant diseases: Status and prospects. In Recent Advances in Biopesticides: Biotechnological Applications; New India Pub.: New Delhi, India; Volume 322.
45. Singh A and Islam M N (2010). In Vitro Evaluation of *Trichoderma* spp against *Phytophthora nicotianae*. Int.J.Expt.Agric.1(1) . 20-25.
46. Sivasithamparam K and Ghisalberti EL (1998). Secondary metabolism in *Trichoderma* and *Gliocladium*. In: *Trichoderma and Gliocladium*, Vol I., (Eds., C.P. Kubicek and G.E.Harman), Taylor and Francis, London. pp.139-191.
47. Smith D and Onions A H S (1994). The preservation and maintenance of living fungi. London:International Mycological Institute.122pp.
48. Van Wees SCM, der Ent S and Pieterse CMJ (2008). Plant immune responses triggered by beneficial microbes. Curr. Opin. Plant Biol. 11, 443–448.
49. Verma M, Satinder KB, Tyagi RD, Surampalli RY and Valero JR (2007). Antagonistic fungi, *Trichoderma* spp. Biochem Engin J. 37: 1-20.
50. Vinale F, Sivasithamparam K, Ghisalberti EL, Marra R, Woo S L and Lorito M (2008). *Trichoderma*–plant–pathogen interactions. Soil Biol. Biochem., 40, 1–10.
51. Viterbo ADA, Chet I and Tas Hyd I (2006). A new hydrophobin gene from the biocontrol agent *Trichoderma asperellum*, is involved in plant root colonization. Mol. Plant Pathol., 7, 249–258.
52. Welkowitz J, Ewen, R. B and Cohen. J (1982). Introductory Statistics for Behavioural Sciences, third ed., Academic Press, New York.

53. Wilson PS., Ahvenniemi PM, Lehtonen MJ, Kukkonen M, Rita Hand Valkonen JPT. (2008). Biological and chemical control and their combined use to control different stages of the *Rhizoctonia* disease complex on potato through the growing season. Ann. Appl. Biol. 153, 307– 320. doi:10.1111/j.17447348.2008.00292.x
54. Zhang J, Chen GY, Li XZ, Hu M, Wang BY, Ruan BH, Zhou H, Zhao LX, Zhou J and Ding ZT (2017). Phytotoxic, antibacterial, and antioxidant activities of mycotoxins and other metabolites from *Trichoderma spp.* Nat. Prod. Res. 31, 2745–2752.

تأثير أربعة أنواع من فطر *Trichoderma spp* على نمو وتطور *Rhizoctonia solani* المعزولة من البطاطس (صنف سبونت) المصابة بمرض القشرة السوداء.

منى نوري عكريم الطاهر أحمد أبو حليقة الصادق محمد غزالة
عبير مسعود القبلاوي أمينة عبدالسلام القمودي

جامعة طرابلس- كلية الزراعة

الملخص باللغة العربية

تم الحصول على عزلات *Rhizoctonia solani* من درنات البطاطس المصابة (صنف سبونت). استخدمت أربع عزلات من الترايكوديرما والمسماة *Trichoderma* *T. album*, *T. longibrachiatum*, *T. viridae*, *harazianum* لاختبار تأثيرها التضادي على نمو وتطور عزلات *R. solani*. تم دراسة التداخل والتضاد بتقنية الزراعة المزدوجة والمواد المتطايرة والغير المتطايرة ما بين العزلات الفطرية المضادة وفطر *R. solani*.

جميع عزلات *Trichoderma spp* أظهرت تأثير معنوي على نمو عزلات *R. solani* المختبرة. أنتجت مناطق تثبيط امام مستعمرات *R. solani* ولكن بدرجات مختلفة. عزلة *Trichoderma harazianum* كانت قادرة على النمو فوق ميسيليوم *R. solani* مقارنة بعزلات الترايكوديرما الأخرى المختبرة. جميع المضادات أكبرت تكون الاجسام الحجرية. أدى ذلك الى تقليل النمو الفطري لعزلات *Rhizoctonia solani*. أشارت النتائج الى أن مدى التثبيط بواسطة *T. harazianum* يشجع على استخدام مضادات محتملة ممتازة قادرة على التحكم في الامراضية لفطر *R. solani* على البطاطس.

الكلمات المفتاحية: البطاطس، العوامل الحيوية الفطرية، التضاد، التطفل *Rhizoctonia solani*,